

Review Article

A Review of the Ichthyofauna of Malaysian Peat Swamp Forest

Sule, H. A.^{1,2}, Ismail, A.^{1*} and Amal, M. N. A.¹

¹*Department of Biology, Faculty of Science, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia*

²*Department of Integrated Science, School of Science, Kogi State College of Education, PMB 1033 Ankpa, Nigeria*

ABSTRACT

A review of literature showed that numerous intensive surveys have been carried out on the ichthyofauna of the peat swamp forests (PSFs) of Malaysia. This review aims to provide a checklist of blackwater fish species in Malaysia from available published literature, and address their economical importance, conservation status and problems of PSFs. A total of 198 peat swamp fish species from 32 families have been recorded in Malaysia. From this number, a total of 114 species from 23 families, representing about 40% of the known fish fauna in Peninsular Malaysia, were recorded from north Selangor PSF. Meanwhile, a total of 49 species belonging to 18 families, 13 species from seven families, 58 species belonging to 19 families, and nine species from five families were recorded from the peat swamps of Perak, Johor, Pahang and East Peninsular Malaysia (parts of Pahang and Terengganu), respectively. Meanwhile, 31 species from 12 families and 40 species belonging to 13 families were recorded from Sabah and Sarawak, respectively. Family Cyprinidae has the highest recorded species, followed by Osphronemidae, Bagridae and Siluridae. The IUCN Red List revealed 12 threatened species facing risk of extinction. The importance of conserving PSFs was outlined and suggestions made in line with the objectives of conservation. Findings from literature revealed that Malaysia's PSFs are rich

in fish diversity, contrary to previous belief, and should therefore be conserved and protected to ensure the richness of their fish diversity.

ARTICLE INFO

Article history:

Received: 18 June 2015

Accepted: 27 May 2016

E-mail addresses:

sule.education001@gmail.com (Sule, H. A.),

aismail@upm.edu.my (Ismail, A.),

mnamal@upm.edu.my (Amal, M. N. A.)

* Corresponding author

Keywords: Ichthyofauna, peat swamp forest, conservation, Malaysia

INTRODUCTION

Peat swamp forests (PSFs) are one of the most unusual and harsh ecosystems in the tropical rainforest biome. The PSFs of Peninsular Malaysia are one of the most threatened (illegal logging, irrigation of oil palm and paddy fields), yet one of the most poorly understood biotopes (Ng et al., 1994). These swamps derive their name from their substrate of peat consisting of plant detritus, which gradually release tannins and organic acids into poorly buffered water and contribute to its characteristically low pH (Ng et al., 1994; Yule, 2010; Posa et al., 2011; Wantzen et al., 2011). Peat swamps are typically deficient in oxygen, which is one consequence of plant decay (Beamish et al., 2003). Peat swamps are generally referred as 'black waters', while waters originating from PSFs are highly acidic with pH values ranging from 3.6 to 5.9, tea-coloured when seen against transmitted light, and black when seen en masse via reflected light (Johnson, 1967ab; 1968). Generally, the acidity of black waters is due to the high concentrations of humic acids and other phenolic acids (Goltenboth, 2006; Irvine et al., 2013).

Most of the black waters of Singapore and Peninsular Malaysia recorded by Wyatt-Smith (1959; 1964), Johnson (1967a; 1968), Anderson (1983) and Whitmore (1984; 1988) have been converted to agriculture, industrial states and residential areas (Ng et al., 1994). The peat swamp forests of Johor, which were of staggering vastness in Peninsular Malaysia, are almost completely degraded. Peat swamp forests

of considerable size in Peninsular Malaysia are now restricted to north Selangor, central Terengganu and Pahang (Ng et al., 1994). In Peninsular Malaysia, the fish fauna of the PSF of north Selangor is comparably well studied and recorded.

The unique characteristics of black waters (dark colour, low dissolved oxygen and high acidity) led to an initial misinterpretation of 'habitat inhospitality', and the conclusion that such habitat will sustain very poor faunal diversity. One of the earliest surveys of the fish fauna of Malaysia peat swamps was by Johnson (1967ab; 1968) who recorded only 26 species in black waters, of which only one is stenotopic to black water. However, the first well-recognised documentation of blackwater fish species of the north Selangor peat swamp forest (NSPSF) was done by Davies and Abdullah (1989). Numerous other surveys have also been conducted on the ichthyofauna of Peninsular Malaysia (Shiraishi et al., 1972; Lim et al., 1982; Mizuno & Furtado, 1982; Zakaria-Ismail, 1990; Ng & Lim, 1991; Ng et al., 1992, 1994; IPT-AWB, 1993; Lee & Ng, 1994; Zakaria et al., 1999; Lee, 2001; Beamish et al., 2003; Rezawaty, 2004; Shah et al., 2006; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013).

The objectives of this review are to provide a checklist of black water fish species in Malaysia, assess their economic importance, update the conservation status of the species and address the conservation problems of PSFs. An exhaustive search of published literature was conducted to

collate lists and determine the distribution of the peat swamp fishes from surveys in Malaysian PSF sites (Figure 1). Fish species provided in this review were recorded from the surveys between 1989 and 2015. Reference was made to Kottelat's catalogue of the fishes of southeast Asia (Kottelat, 2013) to verify and update taxonomic revisions of species.

FISH SPECIES RECORDED IN MALAYSIAN PEAT SWAMP FORESTS

Peat swamp forests have unique water quality and as a result, fish species that can survive and breed in the black water environment are limited. Researchers have reported many species of fish in PSFs that are not found in other habitats (Davies & Abdullah, 1989; Ng et al., 1994). A comprehensive list and a summary of the fish species of Malaysia PSFs are given in Tables 1 and 2.

North Selangor Peat Swamp Forest

The NSPSF has been well studied relative to other PSFs in Malaysia. The most extensive surveys of the fish fauna of NSPSF were done by Davies and Abdullah (1989), IPT-AWB (1993), Ng et al. (1994), Beamish et al. (2003), Ahmad et al. (2013), Ismail et al. (2013) and Siow et al. (2013).


Davies and Abdullah (1989), in a survey of freshwater fish of NSPSF in 1989, recorded 42 fish species (actually 41 species, with the exception of *Osteochilus hasselti*, the synonym of *O. vittatus* recorded as another different species) belonging to 11

families (Table 1). A lot of scientific names, from family to species level, have been changed over the past decade. The most recent comprehensive list of such changes is outlined in Kottelat's catalogue of the fishes of Southeast Asia (Kottelat, 2013).

An intensive survey in 1992 by IPT-AWB (1993) recorded 76 fish species belonging to 24 families (23 with exception of Belontiidae, now included in the family Osphronemidae). This is one of the most successful surveys in NSPSF in terms of diversity. Ng et al. (1994) recorded 47 species of fish living in the black waters of NSPSF, and 14 species of the fishes were stenotopic to acidic black waters. Six unique fish species relatively new to science were first recorded at NSPSF in 1989 (Ng et al., 1994).

Studies by Beamish et al. (2003) conducted from 1997 to 1998 recorded 35 fish species from NSPSF belonging to 14 families (actually 13 because Belontiidae and Luciocephalidae which are now included in Osphronemidae, were recorded separately), while Giam et al. (2012) recorded eight blackwater fish species belonging to the six families from NSPSF.

Recently, during the Selangor Scientific and Biodiversity Peatland Expedition of 2013, Ahmad et al. (2013), Ismail et al. (2013) and Siow et al. (2013) recorded 30 species from 13 families, 41 species from 13 families, and 24 species 10 families, respectively. The comprehensive list of fish species recorded from NSPSF is given in Table 1.



Key:

- | | |
|---|---|
| 1- North Selangor peat swamp forest (NSPSF), Selangor | 6- Segama River, Sabah |
| 2- Paya Berial peat swamp forest (PBPSF), north Perak | 7- Maliau basin, Sabah |
| 3- Black water sites, west Johor | 8- Rajang basin, Sarawak |
| 4- Southeast Pahang peat swamp forest (SEPPSF) | 9- Black water sites, Batang Kerang and Sadong, Sarawak |
| 5- Black water sites, part of Pahang and Terengganu | 10- Black water sites, Nanga Merit area, Sarawak |

Figure 1. Location of PSFs and black water sites surveyed

Table 1
 Checklist of black water fish in Malaysia

| Family | Species | IUCN Status | NSPSF | PBPSF | Johor PSFs | SEPPSF | EP PSFs | Sabah PSFs | Sarawak PSFs | References |
|---------------|--|-------------|-------|-------|------------|--------|---------|------------|--------------|--|
| Akysidae | <i>Acrochordonichthys ischnosoma</i> Bleeker, 1858 | NE | x | | | | | | | Davies & Abdullah, 1989 |
| | <i>Acrochordonichthys rogosus</i> (Bleeker, 1846) | NE | x | | | | | | | IPT-AWB, 1993 |
| | <i>Akysis cf. hendricksoni</i> Alfred, 1986 | LC | x | | | | | | | IPT-AWB, 1993 |
| | <i>Parakysis verrucosus</i> Herre, 1940 | NE | x | | | | | | | IPT-AWB, 1993; Ng et al., 1994 |
| Anabantidae | <i>Anabas testudineus</i> (Bloch, 1792) | DD | x | x | x | | | | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Khairul-Adha et al., 2009; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| Anguillidae | <i>Anguilla borneensis</i> Popta, 1924 | VU | | | | | | x | | Martin-Smith & Tan, 1998 |
| Aplocheilidae | <i>Aplocheilus panchax</i> (Hamilton, 1822) | LC | | x | | | | | | Zakaria et al., 1999; Rezawaty, 2004; Shah et al., 2006 |
| Ariidae | <i>Cryptarius truncatus</i> (Valenciennes, 1840) | NE | | | | | | x | | Martin-Smith & Tan, 1998 |
| | <i>Batrachcephalus mino</i> (Hamilton, 1822) | NE | | | | | | x | | Martin-Smith & Tan, 1998 |
| | <i>Cephalocassis borneensis</i> (Bleeker, 1851) | NE | | | | | | x | | Martin-Smith & Tan, 1998 |
| Engraulidae | <i>Setipinna melanochir</i> (Bleeker, 1849) | NE | | | | | | x | | Martin-Smith & Tan, 1998 |

cont'd Table 1

| | | | | | |
|---|---|---|---|---|--|
| <p>Baltoridae</p> <p><i>Homaloptera cf. nebulosa</i> (Alfred, 1969)</p> <p><i>Homaloptera ogilviei</i> (Alfred, 1967)</p> <p><i>Homaloptera zollingeri</i> (Bleeker, 1853)</p> <p><i>Neohomaloptera johorensis</i> (Herre, 1944)</p> | <p>NE</p> <p>DD</p> <p>LC</p> <p>NE</p> | <p>x</p> <p>x</p> <p>x</p> <p>x</p> | <p>x</p> <p></p> <p></p> <p></p> | <p>x</p> <p></p> <p></p> <p></p> | <p>IPT-AWB, 1993; Ahmad et al., 2005</p> <p>IPT-AWB, 1993</p> <p>IPT-AWB, 1993</p> <p>Ng et al., 1994; Ng & Tan, 1999; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013</p> |
| <p>Barbuccidae</p> <p><i>Barbucca diabolica</i> Roberts, 1989</p> | <p>DD</p> | <p></p> | <p>x</p> | <p>x</p> | <p>Ahmad et al., 2005; Giam et al., 2012</p> |
| <p>Bagridae</p> <p><i>Bagrichthys hypselopterus</i> (Bleeker, 1852)</p> <p><i>Bagrichthys macrocanthus</i> (Bleeker, 1854)</p> <p><i>Bagrichthys macropterus</i> (Bleeker, 1854)</p> <p><i>Hemibagrus baramensis</i> (Regan, 1906)</p> <p><i>Hemibagrus capitulum</i> (Popta, 1904)</p> <p><i>Hemibagrus nemurus</i> (Valenciennes, 1840)</p> <p><i>Hemibagrus planiceps</i> (Valenciennes, 1840)</p> <p><i>Hemibagrus sabanus</i> (Inger & Chin, 1959)</p> | <p>NE</p> <p>NE</p> <p>NE</p> <p>NE</p> <p>NE</p> <p>LC</p> <p>NE</p> <p>NE</p> | <p>x</p> <p>x</p> <p></p> <p>x</p> <p></p> <p>x</p> <p></p> <p></p> | <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> | <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> <p></p> | <p>Davies & Abdullah, 1989</p> <p>Siow et al., 2013</p> <p>Hassan et al., 2010</p> <p>Davies & Abdullah, 1989; Martin-Smith & Tan, 1998</p> <p>Ahmad et al., 2005</p> <p>IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Khairul-Adha et al., 2009; Shah et al., 2006; Jimmy et al., 2010; Ismail et al., 2013; Siow et al., 2013</p> <p>Hassan et al., 2010</p> <p>Martin-Smith & Tan, 1998</p> |

cont'd Table 1

| | | | | | | |
|---|----|---|---|---|---|---|
| <i>Hyalobagrus ornatus</i> (Duncker, 1904) | NE | | | x | x | Ng & Kottelat, 1998; Ahmad et al., 2005; Giam et al., 2012 |
| <i>Leiocassis leiakanthus</i> (Weber & de Beaufort, 1912) | NE | x | | | | Davies & Abdullah, 1989 |
| <i>Leiocassis micropogon</i> (Bleeker, 1852) | NE | x | | | | Ng et al., 1994; Beamish et al., 2003 Siow et al., 2013 |
| <i>Leiocassis poeciloptera</i> (Valenciennes, 1840) | NE | x | | | | IPT-AWB, 1993 |
| <i>Leiocassis stenomus</i> (Valenciennes, 1839) | NE | x | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; |
| <i>Mystus bimaculatus</i> (Volz, 1904) | NE | x | | | | Ismail et al., 2013 |
| <i>Mystus nigriceps</i> (Valenciennes, 1840) | NE | x | x | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Shah et al., 2006; Hassan et al., 2010; Ismail et al., 2013; Siow et al., 2013 |
| <i>Pseudomystus robustus</i> (Inger & Chin, 1959) | NE | | | | x | Martin-Smith & Tan, 1998 |
| Chacidae | LC | | | x | | Giam et al., 2012 |
| Chaudhuriidae | | | | | | |
| <i>Bihunichthys monopterooides</i> (Kottelat & Lim, 1994) | NE | x | | | | IPT-AWB, 1993; Ng et al., 1994 |
| <i>Chendol keelini</i> Kottelat & Lim, 1994 | NE | | | | x | Ahmad et al., 2005 |
| <i>Nagaichthys filipes</i> Kottelat & Lim, 1991 | NE | | | x | x | Giam et al., 2012 |

cont'd Table 1

| | | | | | | | |
|-----------|---|----|---|---|---|---|--|
| Channidae | <i>Channa bankanensis</i> (Bleeker, 1853) | NE | x | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Martin-Smith & Tan, 1998; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Channa gachua</i> (Hamilton, 1822) | LC | x | | | | Ng et al., 1994 |
| | <i>Channa lucius</i> (Cuvier, 1831) | LC | x | x | x | x | Davies & Abdullah, 1989; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Ahmad et al., 2005; Parenti & Lim, 2005; Shah et al., 2006; Khairul-Adha et al., 2009; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Channa marulioides</i> (Bleeker, 1851) | LC | | | x | | Ahmad et al., 2005 |
| | <i>Channa melosoma</i> (Bleeker, 1851) | LC | x | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003 |
| | <i>Channa micropeltes</i> (Cuvier, 1831) | LC | x | | x | | Davies & Abdullah, 1989; Ahmad et al., 2005 |
| | <i>Channa striata</i> (Bloch, 1797) | LC | x | x | x | | Davies & Abdullah, 1989; IPT-AWB, 1993; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Ahmad et al., 2013; Siow et al., 2013 |
| | <i>Channa</i> sp.? | - | | | | x | Shah et al., 2006 |
| Cichlidae | * <i>Oreochromis mossambicus</i> (Peters, 1852) | NT | | | | x | Martin-Smith & Tan, 1998 |

cont'd Table 1

| | | | | | | | | | | |
|-----------|---|----|---|--|---|--|--|---|---|---|
| Clariidae | <i>Clarias batrachus</i> (Linnaeus, 1758) | LC | x | | x | | | | x | IPT-AWB, 1993; Khairul-Adha et al., 2009; Siow et al., 2013 |
| | <i>Clarias leiacanthus</i> Bleeker, 1851 | NE | x | | x | | | x | x | Davies & Abdullah, 1989; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Khairul-Adha et al., 2009; Ahmad et al., 2013 |
| | <i>Clarias macrocephalus</i> Günther, 1864 | NT | x | | x | | | x | x | Rezawaty, 2004; Ahmad et al., 2005; Khairul-Adha et al., 2009; Siow et al., 2013 |
| | <i>Clarias meladerma</i> (Bleeker, 1846) | LC | x | | x | | | | | Ng et al., 1994; Ismail et al., 2013 |
| | <i>Clarias nieuhofii</i> (Valenciennes, 1840) | LC | x | | x | | | x | x | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Khairul-Adha et al., 2009; Ismail et al., 2013 |
| | <i>Encheiloctarias baculum</i> Ng & Lim, 1993 | NE | | | | | | | x | Ng & Tan, 2000; Giam et al., 2012 |
| | <i>Encheiloctarias curtisoma</i> Ng & Lim, 1993 | CR | x | | x | | | x | | IPT-AWB, 1993; Ng et al., 1994; Giam et al., 2012 |
| | <i>Encheiloctarias helioides</i> Ng & Lim, 1993 | CR | | | | | | | x | Giam et al., 2012 |
| | <i>Encheiloctarias prolatus</i> Ng & Lim, 1993 | VU | | | | | | | x | Giam et al., 2012 |
| Cobitidae | <i>Acanthopsoides molobrion</i> (Seibert, 1991) | LC | x | | | | | | | IPT-AWB, 1993 |
| | <i>Acantopsis cf. dialuzona</i> (van Hasselt, 1823) | LC | x | | x | | | | | IPT-AWB, 1993; Rezawaty, 2004 |
| | <i>Kottelatlimia katik</i> (Kottelat & Lim, 1992) | NE | | | | | | x | x | Giam et al., 2012 |

cont'd Table 1

| | | | | | | | | |
|---|----|---|---|---|--|---|--|---|
| <i>Kottelatlimia pristes</i> (Roberts, 1989) | NE | x | | | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003 |
| <i>Pangio kuhlii</i> (Valenciennes, 1846) | NE | x | | | | | | IPT-AWB, 1993 |
| <i>Pangio muraeniformis</i> (de Beaufort, 1933) | NE | | x | | | | | Ahmad et al., 2005 |
| <i>Pangio shufordii</i> (Poopa, 1903) | NE | | x | | | | | Ahmad et al., 2005 |
| <i>Lepidocephalichthys tomaculum</i> (Kottelat & Lim, 1992) | NE | x | | | | | | Ng et al., 1994; Ahmad et al., 2013 |
| Cyprinidae | | | | | | | | |
| <i>Boraras maculatus</i> (Duncker, 1904) | LC | | x | | | | | Ahmad et al., 2005 |
| <i>Barbodes banksi</i> (Herre, 1940) | NE | | x | | | | | Ahmad et al., 2005 |
| <i>Barbodes sealei</i> (Herre, 1933) | NE | | | x | | | | Martin-Smith & Tan, 1998; Jimmy et al., 2010 |
| <i>Barbonymus balleroides</i> (Valenciennes, 1842) | NE | | | x | | | | Martin-Smith & Tan, 1998 |
| <i>Barbonymus gonionotus</i> (Bleeker, 1949) | LC | x | | x | | | | Davies & Abdullah, 1989; Rezawaty, 2004; Siow et al., 2013 |
| <i>Barbonymus schwanefeldii</i> (Bleeker, 1854) | LC | x | | x | | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Rezawaty, 2004; Ismail et al., 2013 |
| <i>Brevibora cheeya</i> (Liao & Tan, 2011) | NE | x | | | | | | Ismail et al., 2013 |
| <i>Crossocheilus cobitis</i> (Bleeker, 1854) | NE | | | | | x | | Martin-Smith & Tan, 1998 |
| <i>Crossocheilus</i> cf. <i>oblongus</i> (Kuhl & van Hasselt, 1823) | LC | x | | | | | | IPT-AWB, 1993 |

cont'd Table 1

| | | | | | | |
|--|----|---|---|---|---|--|
| <i>Cyclocheilichthys apogon</i> (Valenciennes, 1842) | LC | x | x | x | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Zakaria et al., 1999; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Ismail et al., 2013 |
| <i>Cyclocheilichthys heteronema</i> (Bleeker, 1853) | LC | x | | | x | IPT-AWB, 1993; Hassan et al., 2010 |
| <i>Cyclocheilichthys repasson</i> (Bleeker, 1853) | LC | x | | | x | IPT-AWB, 1993; Martin-Smith & Tan, 1998 |
| <i>Desmopuntius hexazona</i> (Weber & de Beaufort, 1912) | NE | x | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Ahmad et al., 2013; Ismail et al., 2013 |
| <i>Desmopuntius johorensis</i> (Duncker, 1904) | NE | x | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Kottelat, 1996; Beamish et al., 2003; Ahmad et al., 2005; Parenti & Lim, 2005; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| <i>Desmopuntius pentazona</i> (Boulenger, 1894) | NE | x | | | x | Parenti & Lim, 2005 |
| <i>Hampala macrolepidota</i> (Kuhl & van Hasselt, 1823) | LC | x | | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Ismail et al., 2013 |
| <i>Hampala sabana</i> Inger & Chin, 1962 | NE | | | | x | Martin-Smith & Tan, 1998 |
| <i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844) | NE | | x | | | Rezawaty, 2004 |

cont'd Table 1

| | | | | | | |
|--|----|---|---|---|---|---|
| <i>Kottelatia brittani</i> (Axelrod, 1976) | NE | x | | | | Zakaria et al., 1999; Rezawaty, 2004 |
| <i>Lobocheilos bo</i> (Popta, 1904) | NE | | x | | | Martin-Smith & Tan, 1998; Hassan et al., 2010 |
| ** <i>Labiobarbus leptocheilus</i> (Valenciennes, 1842) | LC | x | | x | | Davies & Abdullah, 1989; Rezawaty, 2004; Shah et al., 2006 |
| <i>Labiobarbus sabanus</i> (Inger & Chin, 1962) | NE | | | x | | Martin-Smith & Tan, 1998 |
| <i>Luciosoma pellegrinii</i> Popta, 1905 | NE | | | x | | Martin-Smith & Tan, 1998 |
| <i>Macrochirichthys macrochirus</i> (Valenciennes, 1844) | NT | x | | | | Davies & Abdullah, 1989 |
| <i>Mystacoleucus marginatus</i> (Valenciennes, 1842) | LC | x | | | | Davies & Abdullah, 1989; IPT-AWB, 1993 |
| <i>Nematabramis everetti</i> Boulenger, 1894 | NE | | | x | | Martin-Smith & Tan, 1998; Sade & Biun, 2012 |
| <i>Osteochilus chini</i> Karnasuta, 1993 | NE | | | x | | Martin-Smith & Tan, 1998 |
| <i>Osteochilus enneaporos</i> (Bleeker, 1852) | LC | x | | | | IPT-AWB, 1993 |
| <i>Osteochilus ingeri</i> Karnasuta, 1993 | NE | | | x | | Martin-Smith & Tan, 1998 |
| <i>Osteochilus microcephalus</i> (Valenciennes, 1842) | LC | | | x | | Sade & Biun, 2012 |
| <i>Osteochilus spilurus</i> (Bleeker, 1851) | LC | x | x | x | x | Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Giam et al., 2012 |
| <i>Osteochilus vittatus</i> (Valenciennes, 1842) | LC | x | | x | | Davies & Abdullah, 1989; IPT-AWB, 1993; Zakaria et al., 1999; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006 |

cont'd Table 1

| | | | | | | | |
|---|----|---|---|---|--|---|--|
| <i>Osteochilus enneaporos</i> (Bleeker, 1852) | LC | x | | | | x | IPT-AWB, 1993 |
| <i>Oxygaster anomalura</i> (van Hasselt, 1829) | LC | x | x | | | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Shah et al., 2006; Khairul-Adha et al., 2009 |
| <i>Paedocypris micromegethes</i> Kottelat, Britz, Tan & Witte, 2006 | NE | | | | | x | Giam et al., 2012 |
| <i>Paedocypris</i> sp. "North Selangor" | - | x | | | | | Giam et al., 2012 |
| <i>Paedocypris</i> sp. "Pahang" | - | | | x | | | Giam et al., 2012 |
| <i>Parachela hypophthalmus</i> (Bleeker, 1860) | LC | | | x | | | Ahmad et al., 2005 |
| <i>Parachela oxygastroides</i> (Bleeker, 1852) | LC | x | | | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Ismail et al., 2013 |
| <i>Puntioplites bulu</i> (Bleeker, 1851) | DD | x | | | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Martin-Smith & Tan, 1998 |
| <i>Puntius binotatus</i> (Valenciennes, 1842) | LC | x | x | | | | Davies & Abdullah, 1989; Rezawaty, 2004; Shah et al., 2006 |
| <i>Puntius fasciatus</i> (Jerdon, 1849) | LC | x | | | | | Davies & Abdullah, 1989 |
| <i>Puntigrus partipentazona</i> (Fowler, 1934) | LC | | x | | | | Ahmad et al., 2005 |
| <i>Rasbora argyrotaenia</i> (Bleeker, 1849) | NE | | | | | x | Martin-Smith & Tan, 1998 |
| <i>Rasbora bankanensis</i> (Bleeker, 1853) | NE | x | | | | x | IPT-AWB, 1993; Ahmad et al., 2005 |
| <i>Rasbora borapetensis</i> Smith, 1934 | LC | | x | | | | Rezawaty, 2004 |

cont'd Table 1

| | | | | | | |
|--|----|---|---|---|---|---|
| <i>Rasbora cf. sumatrana</i> (Bleeker, 1852) | NE | x | x | x | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Zakaria et al., 1999; Rezawaty, 2004; Shah et al., 2006; Sade & Biun, 2012; Siow et al., 2013 |
| <i>Rasbora dorsiocellata</i> (Duncker, 1904) | NE | x | | | x | Ng et al., 1994; Ahmad et al., 2005 |
| <i>Rasbora maculata</i> (Duncker, 1904) | NE | x | | | | IPT-AWB, 1993; Ng et al., 1994 |
| <i>Rasbora gracilis</i> (Kottelat, 1991) | NE | x | | | x | IPT-AWB, 1993; Beamish et al., 2003; Ahmad et al., 2005 |
| <i>Rasbora kalochroma</i> (Bleeker, 1851) | NE | x | | | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| <i>Rasbora cephalotaenia</i> (Bleeker, 1852) | NE | x | | x | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Ahmad et al., 2013; Ismail et al., 2013 |
| <i>Rasbora dusonensis</i> (Bleeker, 1850) | NE | x | | | | IPT-AWB, 1993; Ismail et al., 2013 |
| <i>Rasbora einthovenii</i> (Bleeker, 1851) | NE | x | | | x | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |

cont'd Table 1

| | | | | | | | |
|--|----|--|---|---|---|---|--|
| <i>Rasbora elegans</i> Volz, 1903 | LC | | | | x | | Ahmad et al., 2005 |
| <i>Rasbora paucisqualis</i> Ahl, 1935 | LC | | | | x | | Ahmad et al., 2005 |
| <i>Rasbora tornieri</i> Ahl, 1922 | LC | | | | | x | Parenti & Lim, 2005 |
| <i>Rasbora trilineata</i> Steindachner, 1870 | LC | | | x | | | Zakaria et al., 1999; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006 |
| <i>Sirium lineatus</i> (Duncker, 1904) | NE | | x | | | | Zakaria et al., 1999; Rezawaty, 2004; Shah et al., 2006; Ismail et al., 2013 |
| <i>Sundanio margariton</i> Conway, Kottelat & Tan, 2011 | NE | | | | | x | Giam et al., 2012 |
| <i>Thynnichthys thynnoides</i> (Valenciennes, 1842) | LC | | x | | | | Davies & Abdullah, 1989 |
| <i>Trigonopoma pauciperforatum</i> (Weber & de Beaufort, 1916) | NE | | x | | x | | IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Ahmad et al., 2005; Khairul Adha et al., 2009; Ahmad et al., 2013; Ismail et al., 2013 |
| <i>Trigonopoma gracile</i> (Kottelat, 1991) | NE | | x | | | | Ismail et al., 2013; Siow et al., 2013 |
| <i>Trigonostigma heteromorpha</i> (Ducker, 1904) | LC | | x | | x | | IPT-AWB, 1993; Zakaria et al., 1999; Ahmad et al., 2005 |
| <i>Oxyeleotris marmorata</i> (Bleeker, 1852) | NE | | x | | x | | Davies & Abdullah, 1989; IPT-AWB, 1993; Rezawaty, 2004 |
| <i>Gobiid</i> sp. | - | | | | | x | Martin-Smith & Tan, 1998 |
| <i>Pseudobiopsis oligactis</i> (Bleeker, 1875) | NE | | x | | | | IPT-AWB, 1993 |

cont'd Table 1

| | | | | | | | | | |
|-----------------|---|----|---|---|---|---|---|---|--|
| Helostomatidae | <i>Helostoma temminckii</i> (Cuvier, 1829) | LC | x | x | x | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Khairul-Adha et al., 2009; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| Mastacembelidae | <i>Macrogynathus aculeatus</i> (Bloch, 1787) | NE | x | | | | | | IPT-AWB, 1993 |
| | <i>Macrogynathus circumcinctus</i> (Hora, 1924) | LC | x | | | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Mastacembelus armatus</i> (La Cèpède, 1800) | LC | | x | | | | | Shah et al., 2006 |
| Nandidae | <i>Nandus nebulosus</i> (Gray, 1835) | LC | x | x | x | | | | IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013 |
| Nemacheilidae | <i>Nemacheilus selangoricus</i> (Duncker, 1904) | DD | x | x | | | | x | Ahmad et al., 2005; Zakaria et al., 1999; Beamish et al., 2003; Ismail et al., 2013 |
| Notopteridae | <i>Chitala chitala</i> (Hamilton, 1822) | NT | x | | | | | | Davies & Abdullah, 1989 |
| | <i>Chitala lopis</i> (Bleeker, 1851) | LC | x | x | | | | x | IPT-AWB, 1993; Ahmad et al., 2005; Shah et al., 2006 |
| | <i>Notopterus notopterus</i> (Pallas, 1976) | LC | x | x | | | | x | Davies & Abdullah, 1989; Rezawaty, 2004; Shah et al., 2006; Ismail et al., 2013 |

cont'd Table 1

| | | | | | | |
|---------------|---|----|---|---|---|---|
| Osphronemidae | <i>Belontia hasselti</i> (Cuvier, 1831) | NE | x | x | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta akarensis</i> Regan, 1910 | NE | | | x | Parenti & Lim, 2005 |
| | <i>Betta bellica</i> (Sauvage, 1884) | LC | x | | x | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta brownorum</i> Witte & Schmidt, 1992 | NE | | | x | Giam et al., 2012 |
| | <i>Betta hipposideros</i> Ng & Kottelat, 1994 | VU | x | | | Ng et al., 1994; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta ibanorum</i> Tan & Ng, 2004 | NE | | | x | Giam et al., 2012 |
| | <i>Betta livida</i> (Ng & Kottelat, 1992) | EN | x | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta imbellis</i> (Ladiges, 1975) | LC | x | | | IPT-AWB, 1993 |
| | <i>Betta persephone</i> Schaller, 1986 | CR | | | x | Giam et al., 2012 |
| | <i>Betta pugnax</i> (Cantor, 1849) | NE | x | x | | Davies & Abdullah, 1989; Zakaria et al., 1999; Beamish et al., 2003; Ismail et al., 2013; Siow et al., 2013 |
| | <i>Betta pulchra</i> Tan & Tan, 1996 | NE | | | x | Giam et al., 2012 |
| | <i>Betta splendens</i> Regan, 1910 | VU | | x | | Rezawaty, 2004; Shah et al., 2006 |

cont'd Table 1

| | | | | | | |
|--|----|---|---|---|---|--|
| <i>Betta taeniata</i> Regan, 1910 | NE | x | | | | Rezawaty, 2004 |
| <i>Betta tomi</i> Ng & Kottelat, 1994 | VU | | x | | | Giam et al., 2012 |
| <i>Betta tussya</i> Schaller, 1985 | NE | | | x | x | Ahmad et al., 2005; Giam et al., 2012 |
| <i>Betta waseri</i> (Krummenacher, 1987) | NE | x | | x | x | IPT-AWB, 1993; Ahmad et al., 2005; Giam et al., 2012 |
| <i>Luciocephalus pulcher</i> (Gray, 1830) | NE | x | | x | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Ahmad et al., 2005; Parenti & Lim, 2005; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| <i>Osphronemus goramy</i> La Cèpède, 1801 | LC | | x | | | Rezawaty, 2004 |
| <i>Parosphromenus alfredi</i> Kottelat & Ng, 2005 | NE | | | | x | Giam et al., 2012 |
| <i>Parosphromenus allani</i> Brown, 1987 | NE | | | | x | Giam et al., 2012 |
| <i>Parosphromenus filamentosus</i> Vierke, 1981 | NE | | | | x | Zakaria et al., 1999 |
| <i>Parosphromenus harveyi</i> (Brown, 1987) | EN | x | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013 |
| <i>Parosphromenus nagy</i> Schaller, 1985 | NE | | | | x | Ahmad et al., 2005; Giam et al., 2012 |
| <i>Parosphromenus tweediei</i> Kottelat & Ng, 2005 | NE | | | | x | Giam et al., 2012 |

cont'd Table 1

| | | | | | |
|--|----|---|---|---|--|
| <i>Sphaerichthys ospromenoides</i> (Canestrini, 1860) | NE | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| <i>Trichopodus leerii</i> (Bleeker, 1852) | NT | x | | x | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |
| <i>Trichopodus pectoralis</i> (Regan, 1910) | LC | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Rezawaty, 2004; Shah et al., 2006; Khairul-Adha et al., 2009 |
| <i>Trichopodus trichopterus</i> (Pallas, 1770) | LC | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Martin-Smith & Tan, 1998; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Khairul-Adha et al., 2009; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| <i>Trichopsis vittata</i> (Cuvier, 1831) | LC | x | | x | IPT-AWB, 1993; Ng et al., 1994; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| Pangasiidae | | | | | |
| <i>Pangasius lithostoma</i> Roberts, 1989 | NE | | | x | Hassan et al., 2010 |
| <i>Pseudolais micronemus</i> (Bleeker, 1846) | DD | | x | | Martin-Smith & Tan, 1998; Ahmad et al., 2005 |

cont'd Table 1

| | | | | | | | | | | | | |
|-----------------|--|----|---|---|---|---|--|--|--|---|---|--|
| Pristolepididae | <i>Pristolepis grootii</i> (Bleeker 1852) | NE | x | | | | | | | | | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ismail et al., 2013 |
| | <i>Pristolepis fasciata</i> (Bleeker, 1851) | LC | x | x | x | x | | | | | | Rezawaty, 2004; Ahmad et al., 2005; Shah et al., 2006; Siow et al., 2013 |
| Siluridae | <i>Kryptopterus bicirrhus</i> (Valenciennes, 1839) | LC | x | | | | | | | | | Davies & Abdullah, 1989; IPT-AWB, 1993 |
| | <i>Kryptopterus limpok</i> (Bleeker, 1852) | NE | | | | | | | | | x | Hassan et al., 2010 |
| | <i>Kryptopterus macrocephalus</i> (Bleeker, 1858) | LC | x | | | | | | | | | Davies & Abdullah, 1989; IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Kryptopterus</i> sp.? | - | | | x | | | | | | | Siow et al., 2013 |
| | <i>Ompok bimaculatus</i> (Bloch, 1794) | NT | x | | x | | | | | | | Davies & Abdullah, 1989; Shah et al., 2006 |
| | <i>Ompok fumidus</i> (Tan & Ng, 1996) | VU | x | | | | | | | | | Beamish et al., 2003; Siow et al., 2013 |
| | <i>Ompok leiacanthus</i> (Bleeker, 1853) | DD | x | | | | | | | | | IPT-AWB, 1993; Ng et al., 1994 |
| | <i>Ompok sabanus</i> Inger & Chin, 1959 | NE | | | | | | | | x | | Martin-Smith & Tan, 1998 |
| | <i>Ompok</i> sp.? | - | | | | | | | | x | | Zakaria et al., 1999; Shah et al., 2006 |
| | <i>Phalacrotonotus apogon</i> (Bleeker, 1851) | LC | | | | | | | | | x | Ahmad et al., 2005 |
| | <i>Phalacrotonotus parvanalis</i> (Inger & Chin, 1959) | NE | | | | | | | | | x | Martin-Smith & Tan, 1998 |

cont'd Table 1

| | | | | | | |
|------------------|---|----|---|---|---|--|
| | <i>Silurichthys hasseltii</i> (Bleeker, 1858) | NE | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Ahmad et al., 2005 |
| | <i>Silurichthys indragiriensis</i> (Volz, 1904) | NE | x | | | Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Silurichthys phaiosoma</i> (Bleeker, 1851) | NE | | | x | Giam et al., 2012 |
| | <i>Wallago leeri</i> Bleeker, 1851 | NE | x | x | | Ahmad et al., 2005; Siow et al., 2013 |
| | <i>Wallago maculatus</i> Inger & Chin, 1859 | NE | | | x | Martin-Smith & Tan, 1998 |
| Symbranchidae | <i>Monopterus albus</i> (Zuiew, 1793) | LC | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Ahmad et al., 2005; Shah et al., 2006; Siow et al., 2013 |
| Syngnathidae | <i>Doryichthys deokhatoides</i> (Bleeker, 1854) | NE | x | | | IPT-AWB, 1993 |
| | <i>Doryichthys martensii</i> (Peters, 168) | NE | x | | | IPT-AWB, 1993 |
| Tetraodontidae | <i>Pao leirus</i> (Bleeker, 1850) | NE | x | | | Davies & Abdulllah, 1989 |
| Toxotidae | <i>Toxotes chatareus</i> (Hamilton, 1822) | NE | | | x | Martin-Smith & Tan, 1998 |
| Zenarchopteridae | <i>Dermogenys pusilla</i> Kuhl & van Hasselt, 1823 | NE | | x | | Shah et al., 2006 |
| | <i>Hemirhamphodon byssus</i> Tan & Lim, 2013 | NE | | | x | Hassan et al., 2010 |
| | <i>Hemirhamphodon kapuasensis</i> Collett, 1991 | NE | | | x | Hassan et al., 2010 |
| | <i>Hemirhamphodon kuekenhali</i> Steindachner, 1901 | NE | | | x | Hassan et al., 2010 |
| | <i>Hemirhamphodon phaiosoma</i> (Bleeker, 1852) | NE | | | x | Hassan et al., 2010 |

cont'd Table 1

| | | | | | | |
|--|----|---|---|---|---|---|
| <i>Hemirhamphodon pogonognathus</i> (Bleeker, 1853) | LC | x | x | x | x | IPT-AWB, 1993; Ng et al., 1994; Zakaria et al., 1999; Beamish et al., 2003; Rezawaty, 2004; Ahmad et al., 2005; Ahmad et al., 2013; Ismail et al., 2013 |
|--|----|---|---|---|---|---|

* Introduced species

**Confusing taxonomy

IUCN Status source: IUCN, (2015)

NE = Not Evaluated, DD = Data Deficient, LC = Least Concern, NT = Near Threatened, VU = Vulnerable, EN = Endangered, CR = Critically Endangered, EW = Extinct in the Wild, EX = Extinct

NSPSF = North Selangor peat swamp forest, PBPSF = Paya Beriah peat swamp forest, SEPPSF = Southeast Pahang peat swamp forest, EP = East Peninsular Malaysia (part of Pahang and Terengganu), PSF's = Peat Swamp Forests

Table 2
Summary of black water fish species recorded in Malaysia

| Peat swamp sites | Location | Region | No. of families | No. of species | Reference |
|---|-----------------------|---------------------|-----------------|----------------|--|
| NSPSF | Selangor | Peninsular Malaysia | 23 | 114 | Davies & Abdullah, 1989; IPT-AWB, 1993; Beamish et al., 2003; Ahmad et al., 2013; Ismail et al., 2013; Siow et al., 2013 |
| PBPSF | Perak | Peninsular Malaysia | 18 | 49 | Zakaria et al., 1999; Rezawaty, 2004; Shah et al., 2006 |
| Black water ditches and ponds of west Johor and Ambat | Johor | Peninsular Malaysia | 7 | 13 | Kottelat, 1996; Ng & Kottelat, 1998; Giam et al., 2012 |
| SEPPSF | Pahang | Peninsular Malaysia | 19 | 58 | Ahmad et al., 2005 |
| Endau drainage, black water sites including ponds and ditches | Pahang and Terengganu | Peninsular Malaysia | 5 | 9 | Kottelat, 1996; Ng & Tan, 1999; Giam et al., 2012 |
| Segama River, Maliau Basin | Sabah | Malaysia Borneo | 12 | 31 | Martin-Smith & Tan, 1998; Sade & Biun, 2012 |
| Rajang Basin, Sadong, Batang Kerang, Nanga Merit | Sarawak | Malaysia Borneo | 13 | 40 | Parenti & Lim, 2005; Khairul-Adha et al., 2009; Hassan et al., 2010; Giam et al., 2012; Tan & Lim, 2013 |

Paya Beriah Peat Swamp Forest, North Perak

The Paya Beriah peat swamp forest (PBPSF), which is located near Bukit Merah reservoir, Perak, has a total area of 5,500 hectares that are bordered by dykes, roads and railway tracks. Like most PSFs in Malaysia, it has been significantly impacted through conversion to residential, industrial and agricultural uses (Ismail & Ali, 2002). Nonetheless, the fish population and its biodiversity in PBPSF are not well documented, in comparison with NSPSF.

The most recognised and well-documented surveys in PBPSF were done by Zakaria et al. (1999), Rezawaty (2004) and Shah et al. (2006).

Zakaria et al. (1999) conducted a study of the swamp-riverine fish populations of two spatially isolated fresh water swamp ecosystems: Beriah Kiri River, which was mistakenly identified as Beriah Kanan River (Shah et al., 2006), located in northern Peninsular Malaysia, and the Ulu Sedili River in southern Peninsular Malaysia. The Beriah Kiri and Ulu Sedili river systems are separated by an extensive mountain

range and a north-south distance of about 900 km, creating an effective dispersal and/or migration barrier (Johnson, 1967a; Prentice & Parish, 1992; Krebs, 2009). Zakaria et al. (1999) reported a total of 24 fish species identified from their study, of which 20 species were from Beriah Kiri and 10 species from Ulu Sedili swamp-riverine area. Eight families were represented from the Beriah Kiri swamp-riverine area. However, only the fish species recorded from the Beriah Kiri swamp-riverine are outlined in this review.

A study from Rezawaty (2004) reported a total of 30 fish species from 12 families in a study carried out on the entire Sungai Beriah drainage system, while during an intensive survey of the PBPSF, specifically in Beriah Kanan, Beriah Kiri and Beriah rivers, Shah et al. (2006) recorded a total of 32 fish species belonging to 13 families (Table 1).

Fishes Recorded from the Peat Swamps of Johor

Information on the ichthyofauna of PSFs of the state of Johor is relatively patchy compared to the NSPSF. Kottelat (1996) and Ng and Kottelat (1998) recorded *Systemus lineatus* and *Hyalobagrus ornatus* respectively from PSF of Johor. Giam et al. (2012) recorded 13 fish species belonging to seven families from black water ditches, streams, pools, remnants of PSF and PSFs of West Johor and Ambat (Table 1).

Fishes Recorded from Southeast Pahang Peat Swamp Forest

Ahmad et al. (2005) recorded 58 fish species (with an additional unidentified species belonging to the family Mastacembelidae) from 17 families (19 following the recent taxonomic revisions detailed in Kottelat's catalogue) during a survey in Bebar River, a large river that flows out of the southeast Pahang peat swamp forest (SEPPSF, which is also known as Pekan peat swamp forest). This survey, which was conducted along Bebar River and Serai River (a tributary of Bebar River), brought the total fish species known in SEPPSF to 65 species (see Ahmad et al., 2005) (see Table 1).

Fishes Recorded from the Peat Swamps of East Peninsular Malaysia

Kottelat (1996) and Ng and Tan (1999) recorded *S. lineatus* and *Neohomaloptera johorensis* respectively from east Peninsular Malaysia. Similarly, Giam et al. (2012) recorded nine fish species from six families from the black waters of east Peninsular Malaysia comprising of part of Malaysia's Johor, Pahang and Terengganu (Table 1).

Fish Species Recorded in the Peat Swamp Forests of Sabah

Sabah, together with Sarawak, Kalimantan Barat, Kalimantan Timur and Brunei is situated in Borneo, the world's third largest island with some 743,107 km² of land area. As a result of "...difficulties of access to the interior part of tropical rainforest, a

lack of reliable dating of igneous rocks, poorly fossiliferous sedimentary rocks and an absence of a coherent stratigraphic scheme for many parts of the island” (Hall & Nichols, 2002), there is only fragmentary information on the geology and fish fauna of the island (Parenti & Lim, 2005). The fish species of Sabah are relatively less surveyed and documented than those of Peninsular Malaysia. Even when intensive surveys are carried out, they have not been widely published by the researchers.

Martin-Smith and Tan (1998) carried out an intensive collection of freshwater fishes over a period of two years from the catchment of the upper Segama River near Danum Valley Field Centre, headwater streams in the catchment of the Kuamut River and from the lower Segama River. They reported a total of 65 species from 20 families. A total of about 30 fish species were recorded in blackwater ditches, ponds and streams from the lower Segama River (Martin-Smith & Tan, 1998).

Also, Sade and Biun (2012) studied the ichthyofauna of Maliau Basin, a saucer-shaped depression enclosed by a mountainous rim in the remote part of Sabah, with an undisturbed flora and fauna. Although 15 fish species were recorded from the study, only three species were recorded from acidic tea-coloured blackwaters in the zone (Table 1).

Fish Species Recorded in the Peat Swamp Forests of Sarawak

Like Sabah, the ichthyofauna of the peat swamps of Sarawak has received little attention in comparison to that of Peninsular

Malaysia. The ichthyofauna of Sarawak is generally neglected in comparison to the other political divisions of Borneo for which detailed checklists have been documented (Parenti & Lim, 2005).

Parenti and Lim (2005) carried out a study of the ichthyofauna of the Rajang Basin in Sarawak, Malaysia, Borneo, and presented a checklist of 164 fish species recorded from the headwaters all the way to the brackish waters, including Belaga and the Balui River, Kapit and the Baleh River, and Sibuan area. A total of seven blackwater fish species were recorded from blackwater ditches and pools along Teku River and remnant of PSF behind the old Sibuan airport.

A previous study from Khairul-Adha et al. (2009) recorded 36 species of fish belonging to 13 families from brownish and blackish water habitats of Batang Kerang in Balai Ringin, Sarawak, during wet and relatively dry seasons. Thirty-two species from 12 families were recorded in the brownish water, while only 12 species from seven families were recorded in the black water habitat. In a study by Hassan et al. (2010), seven out of 15 fish species recorded from 11 families in Nanga Merit area were from a water body with peat-like characteristics.

Recently, Tan and Lim (2013) recorded four species of fish of the genus *Hemirhamphodon* from blackwater ditches and ponds in Sarawak, Malaysia, Borneo, while Giam et al. (2012) recorded 13 fish species belonging to five families from the peat swamps of Rajang and Sadong in Sarawak (Table 1).

TAXONOMIC NOTES

Some of the taxonomic names of the fish species outlined in Table 1 and their placement within families are slightly or completely different from the originally recorded names from the surveys, as a result of the recent changes in the taxonomy of the fish species. A comprehensive list of taxonomic changes is given in Kottelat's catalogue (Kottelat, 2013). Davies and Abdullah (1989) recorded *O. vittatus*, along with its synonym *O. hasselti*, as two completely different species. The species *Belontia hasselti* now classified under the family Osphronemidae was placed in the Anabantidae family. Members of Osphronemidae family were placed in the family Belontia by IPT-AWB (1993) and Ng et al. (1994). In the same vein, *Luciocephalus pulcher* was classified under the family Luciocephalidae. Following the work of Britz (1994), and Kottelat and Whitten (1996), the earlier families of Osphronemidae, Belontiidae and Luciocephalidae were constituted as a single family, Osphronemidae. Recently, Beamish et al. (2003) classified *Nemacheilus selangoricus* under the family Nemacheilidae in the family Bagridae, along with *N. johorensis*. In the same vein, Ahmad et al. (2005) classified *N. selangoricus* of the family Nemacheilidae and *Barbucca diabolica* of the family Barbuccidae under the family Balitoridae. Meanwhile, *Hemirhamphodon progonognathus* of the family Zenarchopteridae was placed in the family Hemiramphidae by IPT-AWB (1993), Ng et al. (1994), Beamish et al. (2003)

and Ahmad et al. (2005). The *Parakysis verucosus* of the family Akysidae was also classified under Parakysidae by IPT-AWB (1993) and Ng et al. (1994), while Siow et al. (2013) recorded *Pristolepis fasciata* of the family Pristolepididae as belonging to the family Nandidae. Also, a recorded fish species belonging to the genus *Kryptopterus* was not designated a species name and the reason was not stated appropriately. Considering there are about 17 distinct species in the genus *Krptopterus* (Kottelat, 2013), it is possible that the sample could be a newly discovered species.

The species recorded as *Puntius lineatus* is now named *Striuntius lineatus*, which is a "preferred" name or "authority", the latter being a synonym. Another species recorded as *Labiobarbus lineatus*, which is excluded from Kottelat's catalogue was regarded as a preferred name for the species *Dangila lineata* in Fish Base, Species 200 and ITIS Catalogue of Life: April 2013, and the NCBI Taxonomy (Torres, 2000; EOL, 2015). *Dangila lineata* was also recorded as a synonym of *L. leptocheilus* in Kottelat's catalogue (Kottelat, 2013), suggesting that both names may actually be referring to the same species. However, Rainboth et al. (2012) considered that *L. lineatus* is a distinct species. There has been some confusion about the taxonomy of the species *L. leptocheilus* (*L. lineatus*). In the taxonomic outline provided in this review, the species is presented as *L. leptocheilus* in line with Kottelat's catalogue. Also, the fish earlier recorded as *Mystus micracanthus* is actually *M. nigriceps* (Roberts, 1993; Ng, 2002; Kottelat, 2013).

ECONOMIC IMPORTANCE OF PEAT SWAMP FISHES

Black water peat swamps are important catchment areas. The peat has great water retention ability and serves as reservoir of rain water, which is utilised in agriculture for irrigating the rice fields adjacent the NSPSF (Low & Balamurugan, 1989). The fish of NSPSF are harvested for consumption and also for the aquarium trade due to their ornamental value (Ng et al., 1994; Ismail et al., 2013) (Table 4). Most of the species harvested for consumption by the local people are sold at very low prices relative to their actual worth (Ng et al., 1994).

A breeding study has been carried out on some of the ornamental fish of peat black water such as pearl gourami (*Trichogaster leeri*), chocolate gourami (*Sphaerichthys osphromenoides*), clown rasbora (*Rasbora kalochroma*), six-banded barb (*P. johorensis*), bellicose Betta (*Betta bellica*), giant fighting fish (*B. waseri*) and pygmy rasbora (*R. maculata*), with the aim of conserving the species and improving their production (FFRC, 1995).

CONSERVATION STATUS OF PEAT SWAMP FISH SPECIES IN MALAYSIA

The conservation status of a species indicates if it still exists and how likely it is to become extinct in the near future (InfoNatura, 2007). The IUCN Red List of Threatened Species is the best known conservation status listing system in the world, which classifies species into nine categories based on the rate of decline of the population, population size, geographical range, and degree of population

and distribution fragmentation (Mace et al., 2008; Biodic, 2013; IUCN, 2014). The categories are “Extinct (EX), Extinct in the Wild (EW), Critically Endangered (CR), Endangered (EN), Vulnerable (VU), Near Threatened (NT), Least Concern (LC), Data Deficient (DD) and Not Evaluated (NE)” (IUCN, 2001; InfoNatura, 2007; IUCN, 2012; IUCN, 2015; IUCN, 2014). A species is EX when there is no reasonable doubt that the last individual has died, as a result of failure to record an individual through exhaustive surveys in known and/or expected habitat, at appropriate times (diurnal, seasonal, annual), throughout its historic range. Extinct in the Wild is used to refer to species that are known only to survive in cultivation, in captivity or as a naturalised population(s) well outside the past range (InfoNatura, 2007; Biodic, 2013; IUCN, 2014).

Out of the 198 black water fish species recorded from Malaysia, 106 (53.54%) species belonging to the category NE indicate that their conservation status has not been evaluated against the criteria. Seven (3.53%) species are DD as a result of inadequate information on their distribution and/or population status. Sixty-two species (31.31%) belonging to the LC category, indicate lowest risk, widespread and abundant species. Six species (3.03%) are NT, signifying likelihood to become endangered in the future. Six (3.03%) and two (1.01%) species are VU and EN, respectively, indicating a highly and very highly risk of extinction in the wild. Besides that, three species (1.52%) are CR and

facing an extremely high risk of extinction in the wild, while six species (3.03%) were not fully classified and so exempted from all the categories (Table 1).

A total of only 12 species (6.70%) are threatened, i.e. under CR, EN and VU category. The IUCN classification may not be a true expression of the conservation status of black water fishes, particularly in Malaysia as so many of the species (57.54%) remain unevaluated against the criteria for classification of threatened species. In data-poor situations, it is not uncommon for listing errors of species that otherwise should be classified as threatened (Gärdenfors, 2000; Keith et al., 2000; Gärdenfors et al., 2001; Keith et al., 2004). For instance, five of the ten black water fish species (*B. brownorum*, *Sundadanio margariton*, *B. ibanorum*, *Parosphromenus allani* and *H. ornatus*), listed as “most VU” by Giam et al., (2012) using the criteria of decline in population, geographical range and basin extinction, were classified under the NE in Red List category (Tables 1 and 3).

It is important to point out that the species recorded as ‘rare’ at a distinct time may be more abundant at other times. For example, the abundance of individual fish and species composition vary during the wet and dry seasons, which may be related to variations in migratory movements of fish species (Ng et al., 1994; Renato et al., 2000; Khairul-Adha et al., 2009). Some species migrate from downstream to upper reaches of a river during high water levels for breeding or food, then migrate back to the lower reaches after spawning as the water level reduces (Lowe-McConnell, 1975;

Welcomme, 1979). Therefore, conservation surveys should be exhaustive, in all known and/or expected habitats, at appropriate times covering seasons of high and low water levels over an extended period of time before a species is pronounced as rare or threatened.

Considering the extensive destruction of tropical PSFs worldwide and the decline in PSF biodiversity, more black water fish species will fall into the threatened (CR, EN and VU) category if evaluated against the criteria. However, it is important to note that the category of threat simply provides an assessment of the extinction risk under current circumstances, and is not necessarily sufficient to determine priorities for conservation action (Mace & Lande, 1991).

CONSERVATION OF PEAT SWAMP FORESTS

Peat swamp forests are an important component of the world’s wetlands, providing a wide variety of goods and services in the form of carbon sequestration, flood mitigation and globally important biodiversity reservoirs (Parish, 2002; UNDP, 2006; Posa et al., 2011; Ismail et al., 2013), with many endemic species (Kottelat & Ng, 2005; Kottelat & Widjanarti, 2005; UNDP, 2006; Tan & Kottelat, 2009; Conway et al., 2011; Conway & Kottelat, 2011; Posa et al., 2011; Giam et al., 2012; Taskforce REDD, 2012; PIU-SERT, 2013; Muchlisin et al., 2015).

Malaysian PSFs make up about 75% of the country’s total wetlands, with 80%

Table 3
Peat swamp fishes listed as Threatened in IUCN Red List

| Family | Species | Status | Reference |
|---------------|---------------------------------|--------|--|
| Anguillidae | <i>Anguilla borneensis</i> | VU | Martin-Smith & Tan, 1998 |
| Clariidae | <i>Encheloclarias curtisoma</i> | CR | IPT-AWB, 1993; Ng et al., 1994; Giam et al., 2012 |
| | <i>Encheloclarias keliodes</i> | CR | Giam et al., 2012 |
| | <i>Encheloclarias prolatus</i> | VU | Giam et al., 2012 |
| Osphronemidae | <i>Ompok fumidus</i> | VU | Beamish et al., 2003; Siow et al., 2013 |
| | <i>Betta hipposideros</i> | VU | Ng et al., 1994; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta livida</i> | EN | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013 |
| | <i>Betta splendens</i> | VU | Rezawaty, 2004; Shah et al., 2006 |
| | <i>Betta tomi</i> | VU | Giam et al., 2012 |
| | <i>Betta persephone</i> | CR | Giam et al., 2012 |
| | <i>Parosphronemus harveyi</i> | EN | IPT-AWB, 1993; Ng et al., 1994; Beamish et al., 2003; Giam et al., 2012; Ahmad et al., 2013; Ismail et al., 2013 |

found in east Malaysia (Sabah 8%; Sarawak 72%) and 20% in Peninsular Malaysia. Peat swamp forests in Malaysia have undergone severe degradation over the years. For example, the PSF cover of NSPSF was estimated to be 0.67 million hectares in 1981 but reduced drastically to 0.34 million hectares in the 1990s (UNDP, 2006) and there has been further destruction since then. As the nation becomes aware of the need to conserve PSFs, some percentage of PSFs have been protected within Permanent Forest Reserves and stateland forests (UNDP, 2006). However, this protection is inadequate. Agricultural conversion and fire have destroyed PSF within these protected areas and furthermore, unless the entire forests are protected, any drainage around the edges impacts the entire forest.

The main threats of PSFs in Malaysia are forestry (overexploitation), illegal logging, pollution (including oil, industrial, nutrient and sedimentation), waste disposal, airports, land reclamation, large scale land conversion for agriculture, industrialisation and settlement-urbanisation (Ng & Shamsudin, 2001; Chong et al., 2010). In addition, draining and clearing of the PSFs for agriculture and palm oil plantations have resulted in peat land fires (Parish, 2002; Page et al., 2011), which contribute to the annually recurring episodes of transboundary haze pollution in the Southeast Asian region (Lo & Parish, 2013).

Several studies on the diversity of fish in PSFs have revealed the existence of economically important fishes for consumption, aquarium trade and

Table 4
Commonly consumed and ornamental black water fish species in Malaysia

| Consume | Ornamental |
|-----------------------------------|--------------------------------------|
| <i>Anabas testudineus</i> | <i>Betta bellica</i> |
| <i>Belontia hasselti</i> | <i>Betta livida</i> |
| <i>Channa bankanensis</i> | <i>Betta hipposideros</i> |
| <i>Channa lucius</i> | <i>Belontia hasselti</i> |
| <i>Channa melosoma</i> | <i>Channa gachua</i> |
| <i>Clarias meladerma</i> | <i>Helostoma temminkii</i> |
| <i>Clarias nieuhofii</i> | <i>Hemirhamphodon progonognathus</i> |
| <i>Clarias leiacanthus</i> | <i>Kryptopterus macrocephalus</i> |
| <i>Helostoma temminkii</i> | <i>Leiocassis micropogon</i> |
| <i>Kryptopterus macrocephalus</i> | <i>Luciocephalus pulcher</i> |
| <i>Monopterus albus</i> | <i>Macrognathus circumcinctus</i> |
| <i>Mystus bimaculatus</i> | <i>Mystus bimaculatus</i> |
| <i>Macrognathus circumcinctus</i> | <i>Osteochilus spilurus</i> |
| <i>Hemibagrus nemurus</i> | <i>Parosphromenus harveyi</i> |
| <i>Pristolepis grootii</i> | <i>Desmopuntius hexazona</i> |
| <i>Trichopodus leerii</i> | <i>Desmopuntius johorensis</i> |
| <i>Trichopodus pectoralis</i> | <i>Rasbora cephalotaenia</i> |
| <i>Trichopodus trichopterus</i> | <i>Rasbora dorsiocellata</i> |
| | <i>Rasbora einthovenii</i> |
| | <i>Rasbora gracilis</i> |
| | <i>Rasbora kalochroma</i> |
| | <i>Rasbora maculata</i> |
| | <i>Trigonopoma pauciperforatum</i> |
| | <i>Silurichthys hasseltii</i> |
| | <i>Sphaerichthys osphromenoides</i> |
| | <i>Trichopodus leerii</i> |
| | <i>Trichopodus trichopterus</i> |

endangered species (IPT-AWB, 1993; Ng et al., 1994; Lee, 2001; Beamish et al., 2003), where the majority of fish are highly dependent on the rivers in the PSFs (Beamish et al., 2003; Yule, 2010). The degradation of PSFs is expected to affect the fish community structure in such a way that extinction of some already endangered species in the near future is probable. Gibson et al. (2011) reported that converted land-

use types vary in their ability to support forest biodiversity, generally having a reduced capacity for biodiversity support than in their original unconverted form. Giam et al. (2012) predicted the number of fish species that would become extinct under different land-use conditions using the Matrix-Calibrated Species-Area Model (MCSAM) (Koh & Ghazoul, 2010) and Mote Carlo simulations to project PSF

basins extinctions. Under the scenario, the 10 most vulnerable species (*Encheloclarias prolatus*, *B. brownorum*, *S. goblinus*, *S. margarition*, *B. ibanorum*, *B. burdigala*, *E. tapeinopterus*, *Paedocypris progenetica*, *P. allani* and *H. ornatus*) were recognised and predicted to be extinct by 2050, if the present rate of PSFs conversion continues.

Meanwhile, Beamish et al. (2003) reported 22 species of fish from 43 sites in the NSPSF in 1998 as against 33 species from 27 sites during the preceding year. The PSF was largely forested during the 1997 survey, but the area was cleared and planted with oil palms in 1998. Thirteen species caught in 1997 were not caught in 1998. This supports the prediction by Giam et al. (2012) the extinction of fish species following PSF land conversion. Generally, riparian vegetation acts as a source of energy and matter (Kindler, 1998), and contributes matter to the PSF ecosystem through production of leaf litter (Tabacchi et al., 1998). Modifications in riparian vegetation through logging affect the structure and processes within the peat swamps. It leads to the alteration of the swamp characteristics, reduction of food resources (Tabacchi et al., 1998; Wright & Flecker, 2004) and subsequent loss of biodiversity (Bruenig & Droste, 1995).

Peat swamp forests also serve as an important global carbon storehouse (Parish, 2002; Chimer & Ewel, 2005; Jauhiainen et al., 2005; Rydin & Jeglum, 2006; UNDP, 2006). The PSFs of Malaysia and Indonesia alone store 67 gigatons of carbon in peat, which represents 75% of total tropical peat

soil carbon storage (Page et al., 2011). Large scale conversion of this carbon sink, as is happening today, is expected to and will severely impact the earth's climate (Couwenberg et al., 2009; van der Werf et al., 2009; Miettinen et al., 2012) unless drastic measures are put in place to conserve PSFs of the world.

In the discussion of conservation of PSFs fish biodiversity, safe fishing techniques have always been ignored. Researchers, and fishermen alike, employ varying fishing techniques in PSFs and have always been more concerned with increase in catchability than safety and conservation of the fishes. One fishing technique, electro-fishing, that poses a threat to the health of fish has been employed in several surveys in Malaysia PSFs (Martin-Smith & Tan, 1998; Beamish et al., 2003; Shah et al., 2006; Siow et al., 2013). Electro-fishing establishes an electric field in the water (Lamarque, 1990) creating an epileptic response in fish resulting from electric shock to the central nervous system (Reynolds & Kolz, 1993; Sharber & Black, 1999). The reported health effects on fish include internal haemorrhaging and skin discolorations (Dalbey et al., 1996; Kocovsky et al., 1997; Muth & Ruppert, 1997; Thompson et al., 1997; Ainslie et al., 1998; Habera et al., 1999), spinal injuries (Kocovsky et al., 1997), lacticidosis and disturbance of the inter-renal stress response (Mitton & McDonald, 1994), retarded growth (Dalbey et al., 1996; Thompson et al., 1997; Ainslie et al., 1998; Hughes, 1998), and low gamete viability (Muth & Ruppert, 1997; Koupal et al., 1997). Although the

effects of electro-fishing on PSF biodiversity is minor compared to the enormous loss due to PSF drainage, clearance and fire, it is appropriate that only safe fishing techniques such as the use of scoop nets, cast nets, etc. should be employed in a unique habitat such as peat swamps.

CONCLUSION

It is apparent from the ongoing that one of the strongest justifications for conservation of peat swamps is the existing ichthyofauna. Therefore, maintaining fish populations of PSFs is very important as this will require maintaining the natural water tables of the swamps with seasonal flooding. Apart from the necessity to maintain appropriate aquatic habitats for the fish, another important reason is that when peat dries out due to drainage, it is extremely inflammable and thus, becomes highly vulnerable to destruction by fire (Langner et al., 2007; Langner & Siegert, 2009; Page et al., 2009; Posa et al., 2011).

A review of literature on PSF fish and also the IUCN Red List revealed the likelihood of many important fishes becoming extinct in the near future if the present PSFs degradation continues. Moreover, there is also an inevitable risk of increased global warming when the tons of carbon stored in peat lands is released through degradation. Consequently, a concerted effort for conserve regional PSFs is essential, and requires action plans involving all stakeholders (Azmi et al., 2009). The present measures aimed at protecting only some parts of the peat

swamp forests in Malaysia are totally inadequate (and in any case, they are largely ignored).

REFERENCES

- Ahmad, A. B., Ahmad, M. F., & Nek, S. A. R. T. (2013). *Freshwater fishes recorded from north Selangor peat swamp forest, Selangor*. Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest, 28th November 2013, Quality Hotel, Shah Alam, Selangor, Malaysia. pp.72–76.
- Ahmad, A., Mazlan, A. G., Syed Ahmad Rizal, T. N., Abd. Rahman, A. R., Jamil, M., Jacqueline, L., & Abdullah, S. (2005). *Freshwater fishes of Sungai Bebar and adjacent areas*. In Summary Findings PSF Technical Series No. 4. Peat Swamp Forest Project. UNDP, GEF, Pahang Forestry Department and Universiti Kebangsaan. pp.145–148.
- Ainslie, B. J., Post, J. R., & Paul, A. J. (1998). Effects of pulsed and continuous DC electrofishing on juvenile rainbow trout. *North American Journal of Fisheries Management*, 18(4), 905–18.
- Ali, A. (1999). *A case study on the conservation and management of freshwater wetlands in Malaysia*. Proceedings of Wetlands, Awareness, Local People and the Ramsar Convention in the Mekong River Basin. In O. Callaghan (Ed.), *Can Local People Play a Role in the Wise Use of Wetlands* (pp. 22–32). Phnom Penh, Cambodia.
- Anderson, J. A. R. (1983). The tropical peat swamps of western Malesia. In A. J. P. Gore (Ed.), *Systems of the World 4B. Mires: Swamp, Bog, Fen and Moor* (pp. 181–99). Amsterdam: Elsevier Sci. Publ.
- Azmi, M. I., Cullen, R., Bigsby, H., & Awang, N. A. G. (2009). *The existence value of peat swamp forest in Peninsular Malaysia*. Paper presented at New Zealand Agriculture and Resource Economics Society (NZARES) Conference. 27–28 August

2009. Tahuna Beach Resort, Nelson, New Zealand.
- Beamish, F. W. H., Beamish, R. B., & Lim, S. L. (2003). Fish assemblages and habitat in a Malaysian blackwater peat swamp. *Environmental Biology of Fishes*, 68(1), 1–13.
- Biodic. (2013). *Threatened Species*. Retrieved from www.biodic.go.jp.
- Britz, R. (1994). Ontogenetic features of *Luciocephalus* (Perciformes, Anabantoidei) with a revised hypothesis of anabantoid intrarelationships. *Zoological Journal of the Linnaean Society*, 112(4), 491–508.
- Bruenig, E. F., & Droste, H. J. (1995). Structure, dynamics and management of rainforests on nutrient-deficient soils in Sarawak. In R. B. Primack & T. E. Lovejoy (Eds.), *Ecology, Conservation and Management of Southeast Asian Rainforests* (pp. 41–53). Yale University Press.
- Chimer, R. A., & Ewel, K. C. (2005). *A tropical freshwater wetland: II. Production, decomposition and peat formation*. Asean Wetland Bureau/WWF Malaysia. Kuala Lumpur, Malaysia.
- Chong, V. C., Lee, P. K. Y., & Lau, C. M. (2010). Diversity, extinction risk and conservation of Malaysian fishes. *Journal of Fish Biology*, 76(9), 2009–2066.
- Conway, K. W., & Kottelat, M. (2011). *Boraras naevus*, a new species of miniature and sexually dichromatic freshwater fish from Peninsular Thailand (Ostariophysi: Cyprinidae). *Zootaxa*, 3002, 45–51.
- Conway, K. W., Kottelat, M., & Tan, H. H. (2011). Review of the Southeast Asian miniature cyprinid genus *Sundanio* (Ostariophysi: Cyprinidae) with descriptions of seven new species from Indonesia and Malaysia. *Ichthyological Exploration of Freshwaters*, 22(3), 251–288.
- Couwenberg, J., Dommain, R., & Joosten, H. (2009). Greenhouse gas fluxes from tropical peatlands in Southeast Asia. *Global Change Biology*, 16(6), 1715–1732.
- Dalbey, S. R., McMahon, T. E., & Fredenberg, W. (1996). Effect of electrofishing pulse shape and electrofishing-induced spinal injury on long-term growth and survival of wild rainbow trout. *North American Journal of Fisheries Management*, 16(3), 560–569.
- Davies, J., & Abdullah, A. R. (1989). *Freshwater fish survey of the north Selangor peat swamp forest*. Asian Wetland Bureau Publication No. 46, IPT Asian Wetland Bureau/WWF Malaysia, Kuala Lumpur.
- Encyclopedia of Life. (2015). Synonyms of *Labiobarbus lineatus*. *EoL Online*. Retrieved from <http://eol.org/pages/210213/names/synonyms>.
- Freshwater Fisheries Research Centre (FFRC). (1995). Annual report. *FFRC Online*. Retrieved from <http://www.fri.gov.my/pppat/page11-1.html>.
- Gärdenfors, U. (2000). Population viability analysis in the classification of threatened species: Problems and potentials. *Ecological Bulletins*, 48, 181–190.
- Gärdenfors, U., Hilton-Taylor, C., Mace, G. M., & Rodríguez, J. P. (2001). The application of IUCN Red List criteria at regional levels. *Conservation Biology*, 15(5), 1206–1212.
- Giam, X., Koh, L. P., Tan, H. H., Miettinen, J., Tan, H. T. W., & Ng, P. K. L. (2012). Global extinctions of peat swamp fishes follow peatland conversion in Sundaland. *Frontiers in Ecology and the Environment*, 10(9), 465–470.

- Gibson, L., Lee, T. M., & Koh, L. P. (2011). Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature*, 478(7369), 378–381.
- Goltenboth, F. (2006). *Ecology of Insular Southeast Asia: The Indonesian Archipelago*. In F. Goltenboth, K. H., Timotius, P. P., Milan, & J. Margraf (Eds.). Elsevier.
- Habera, J. W., Strange, R. J., & Saxton, A. M. (1999). AC electrofishing injury of large brown trout in low-conductivity streams. *North American Journal of Fisheries Management*, 19(1), 120–126.
- Hall, R. & Nichols, G. (2002). Cenozoic sedimentation and tectonics in Borneo: Climatic influences on orogenesis. In S. J. Jones & L. Frostick (Eds.), *Sediment Flux to Basins: Causes, Controls and Consequences* (pp. 5–22), Special Publications. London: Geological Society.
- Hassan, R., Esa, F. A., & Harith, M. N. (2010). Fish assemblages in Nanga Merit area, Kapit, Sarawak. *The Sarawak Museum Journal*, 88, 274–284.
- Hughes, N. F. (1998). Reduction in growth due to electrofishing and tagging may change interannual movement behavior of stream salmonids: Evidence from Arctic grayling in an interior Alaskan stream. *Transactions of the American Fisheries Society*, 127(6), 1072–1077.
- InfoNatura. (2007). *About the data: Conservation status*. Retrieved from www.natureserve.org.
- IPT-Asian Wetland Bureau. (IPT-AWB) (1993). *Fish sampling demonstration in north Selangor peat swamp forest*. AWB Publication No. 89.
- Irvine, K., Vermette, S., & Mustafa, F. B. (2013). The “black waters” of Malaysia: tracking water quality from the peat swamp forest to the sea. *Sains Malaysiana*, 42(11), 1539–1548.
- Ismail, A., Amal, M. N. A., Abdullah, T., Johari, S., Azizul, A., Nur-Illiani, H., & Amiruddin, M. H. (2013). *Fish diversity in north Selangor peat swamp forest*. Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest, 28th November 2013, Quality Hotel, Shah Alam, Selangor, Malaysia. pp.57–71.
- Ismail, W. R., & Ali, A. (2002). Managing our wetlands: Some results of the function of a wetland in Perak, Malaysia. *Malaysian Journal of Environmental Management*, 3, 83–105.
- IUCN. (2001). *IUCN Red List Categories and Criteria*. Version 3.1. Gland, Switzerland and Cambridge, UK: IUCN Species Survival Commission.
- IUCN. (2012). *IUCN Red List Categories and Criteria*. Version 3.1. Gland, Switzerland and Cambridge, UK: IUCN.
- IUCN. (2014). *Guidelines for using the IUCN Red List Categories and Criteria* (Version 11). IUCN Standards and Petitions Subcommittee.
- IUCN. (2015). *IUCN Red List of Threatened Species*. Retrieved from www.iucnredlist.org.
- Jauhainen, J., Takahashi, H., Heikkinen, J. E. P., Martikainen, P. J., & Vasanderet, H. (2005). Carbon fluxes from a tropical peat swamp forest floor. *Global Change Biology*, 11(10), 1788–1797.
- Johnson, D. S. (1967a). Distributional patterns of Malayan freshwater fish. *Ecological Society of America*, 48(5), 722–730.
- Johnson, D. S. (1967b). On the chemistry of freshwaters in southern Malaya and Singapore. *Archiv für Hydrobiologie*, 63(4), 477–496.
- Johnson, D. S. (1968). Malayan blackwaters. In R. Misra & B. Gopal (Eds.), *Proceedings of the Symposium on Recent Advances in Tropical Ecology* (pp. 303–310). International Society for Tropical Ecology, Varanasi.
- Keith, D. A., Auld, T. D., Ooi, M. K. J., & Mackenzie, B. D. E. (2000). Sensitivity analyses of decision rules in World Conservation Union (IUCN) Red

- List criteria using Australian plants. *Biological Conservation*, 94(3), 311–319.
- Keith, D. A., McCarthy, M. A., Regan, H., Regan, T., Bowles, C., Drill, C., ... & Ruckelshaus, M. (2004). Protocols for listing threatened species can forecast extinction. *Ecology Letters*, 7(11), 1101–1108.
- Khairul-Adha, A. R., Siti, K. D., Siti, S. S., Aziz, A., Yuzine, E., & Eza, R. I. (2009). Freshwater fish diversity and composition in Batang Kerang floodplain, Balai Ringin, Sarawak. *Pertanika Journal of Tropical Agricultural Science*, 32(1), 7–16.
- Kindler, J. (1998). Linking ecological and development objectives: Trade-offs and imperatives. *Ecological Applications*, 8(3), 591–600.
- Kocovsky, P. M., Gowan, C., Fausch, K. D., & Riley, S. C. (1997). Spinal injury rates in three wild trout populations in Colorado after eight years of backpack electrofishing. *North American Journal of Fisheries Management*, 17(2), 308–313.
- Koh, L. P. & Ghazoul, J. (2010). A matrix-calibrated species–area model for predicting biodiversity losses due to land-use change. *Conservation Biology*, 24(4), 994–1001.
- Kottelat, M. (1996). The identity of *Puntius eugrammus* and diagnoses of two new species of striped barb (Teleostei: Cyprinidae) from Southeast Asia. *The Raffles Bulletin of Zoology*, 44(1), 301–316.
- Kottelat, M. (2013). The fishes of the inland waters of southeast Asia: A catalogue and core bibliography of the fishes known to occur in freshwaters, mangroves and estuaries. *The Raffles Bulletin of Zoology, Supplement No. 27*, 1–663.
- Kottelat, M., Britz, R., Tan, H. H., & Witte, K. E. (2006). *Paedocypris*, a new genus of southeast Asian cyprinid fish with a remarkable sexual dimorphism, comprises the world's smallest vertebrate. *Proceedings of the Royal Society B*, 273(1589), 895–899.
- Kottelat, M., & Ng, P. K. L. (2005). Diagnoses of six new species of *Parosphromenus* (Teleostei: Osphronemidae) from Malay Peninsular and Borneo, with notes on other species. *The Raffles Bulletin of Zoology*, 13, 101–113.
- Kottelat, M., & Whitten, A. J. (1996). *Freshwater fishes of western Indonesia and Sulawesi: Additions and corrections*. Hong Kong: Periplus.
- Kottelat, M., & Widjanarti, E. (2005). The fishes of Danau Sentarum National Park and the Kapuas Lakes area, Kalimantan Barat, Indonesia. *The Raffles Bulletin of Zoology*, 13, 139–173.
- Koupal, K. D., Satterfield, J. R., & Flickinger, S. A. (1997). Comparative gear selectivity for male walleyes and influence of method of capture on resultant hatching success. *The Progressive Fish-Culturist*, 59(3), 218–221.
- Krebs, C. J. (2009). *Ecology: The Experimental Analysis of Distribution and Abundance* (6th ed). Benjamin Cummings. San Francisco.
- Lamarque, P. (1990). Fishing With Electricity: Applications in Freshwater Fisheries Management. In I. G. Cowx, & P. Lamarque (eds.), *Fishing News Books*. Oxford, England.
- Langner, A., Miettinen, J., & Siegert, F. (2007). Land cover change 2002–2005 in Borneo and the role of fire derived from MODIS imagery. *Global Change Biology*, 13(11), 2329–2340.
- Langner, A. & Siegert, F. (2009). Spatiotemporal fire occurrence in Borneo over a period of 10 years. *Global Change Biology*, 15(1), 48–62.
- Lee, K. Y. (2001). *Fish community of the north Selangor peat swamp forest*. (M.Sc. thesis). University of Malaya, Kuala Lumpur, Malaya.
- Lee, P. G., & Ng, P. K. L. (1994). The systematics and ecology of snakeheads (Pisces: Channidae) in Peninsular Malaysia and Singapore. *Hydrobiologia*, 285(1-3), 59–74.

- Lim, R. P., Furtado, J. I., & Morley, R. J. (1982). General description of Tasek Bera. In J.I. Furtado, & S. Mori (Eds.), *Tasek Bera* (pp. 7-54). Monographiae Biologicae. Dordrecht: Springer Netherlands.
- Lo, J., & Parish, F. (2013). *Peatlands and climate change in Southeast Asia*. ASEAN Peatland Forests Project and Sustainable Management of Peatland Forests Project.
- Low, K. S., & Balamurugan, G. (1989). *A Preliminary Hydrological Investigation of the North Selangor Peat Swamp Forest*. Asean Wetland Bureau/ WWF Malaysia. Kuala Lumpur.
- Lowe-McConnell, R. H. (1975). *Fish Communities in Tropical Freshwater*. Longham Inc. New York.
- Mace, G. M., Collar, N. J., Gaston, K. J., Hilton-Taylor, C., Akçakaya, H. R., Leader-Williams, N., Milner-Gulland, E. J., & Stuart, S. N. (2008). Quantification of extinction risk: IUCN's system for classifying threatened species. *Conservation Biology*, 22(6), 1424–1442.
- Mace, G. M., & Lande, R. (1991). Assessing extinction threats: Toward a reevaluation of IUCN threatened species categories. *Conservation Biology*, 5(2), 148–157.
- Martin-Smith, M. K. & Tan, H. H. (1998). Diversity of freshwater fishes from eastern Sabah: Annotated checklist for Danum Valley and a consideration of inter- and intra-catchment variability. *The Raffles Bulletin of Zoology*, 46(2), 573–604.
- Miettinen, J., Shi, C., & Liew, S. C. (2012). Two decades of destruction in southeast Asia's peat swamp forests. *Frontiers in Ecology and the Environment*, 10(3), 124–128.
- Mitton, C. J. A., & McDonald, D. G. (1994). Consequences of pulsed DC electrofishing and air exposure to rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences*, 51(8), 1791–1798.
- Mizuno, N., & Furtado, J. I. (1982). Ecological notes on fishes. In J.I. Furtado & S. Mori (Eds.), *Tasek Bera* (pp. 321-354). Dordrecht, Springer Netherlands.
- Muchlisin, Z. A., Akyun, Q., Rizka, S., Fadli, N., Sugianto, S., Halim, A., & Siti-Azizah, M. N. (2015). Ichthyofauna of Tripa peat swamp forest, Aceh province, Indonesia. *Check List*, 11, 1–9.
- Muth, R. T. & Ruppert, J. B. (1997). Effects of electrofishing fields on captive embryos and larvae of razorback sucker. *North American Journal of Fisheries Management*, 17(1), 160–166.
- Ng, H. H. (2002). The identity of *Mystus nigriceps* (Valenciennes in Cuvier & Valenciennes, 1840), with the description of a new bagrid catfish (Teleostei: Siluriformes) from southeast Asia. *The Raffles Bulletin of Zoology*, 50(1), 161–168.
- Ng, H. H. & Kottelat, M. (1998). *Hyalobagrus*, a new genus of miniature bagrid catfish from southeast Asia (Teleostei: Siluriformes). *Ichthyological Exploration of Freshwaters*, 9, 335–346.
- Ng, H. H. & Tan, H. H. (1999). The fishes of Endau Drainage, Peninsular Malaysia with descriptions of two new species of catfishes (Teleostei: Akysidae, Bagridae). *Zoological Studies*, 38(3), 350–366.
- Ng, P. K. L. & Lim, K. K. P. (1991). The identity of *Ophicephalus cyanospilos* Bleeker from Sumatra and a new record of *Channa bankanensis* (Bleeker) from Peninsular Malaysia (Pisces: Channidae). *The Raffles Bulletin of Zoology*, 39, 119–130.
- Ng, P. K. L., Tay, J. B., & Lim, K. K. P. (1994). Diversity and conservation of blackwater fishes in Peninsular Malaysia, particularly in the north Selangor peat swamp forest. *Hydrobiologia*, 285(1-3), 203–218.

- Ng, P. K. L., Tay, J. B., Lim, K. K. P., & Yang, C. M. (1992). *The conservation of the fish and other aquatic fauna of the north Selangor peat swamp forest and adjacent areas*. Kuala Lumpur: AWB Publication No. 81.
- Ng, T. P. & Shamsudin, I. (2001). *Common trees in peat swamp forests of Peninsular Malaysia. Research Pamphlet No. 124*. Forest Research Institute Malaysia (FRIM). Kepong, Selangor.
- Page, S. E., Hoscilo, A., Langner, A., Tansey, K., Siegert, F., Limin, S., & Rieley, J. O. (2009). Tropical peatland fires in southeast Asia. In M.A. Cochrane (Ed.), *Tropical Fire Ecology: Climate Change, Land Use and Ecosystem Dynamics* (pp. 263–287). USA: Springer.
- Page, S. E., Rieley, J. O., & Banks, C. J. (2011). Global and regional importance of the tropical peatland carbon pool. *Global Change Biology*, 17(2), 798–818.
- Parenti, L. R. & Lim, K. K. P. (2005). Fishes of the Rajang Basin, Sarawak, Malaysia. *The Raffles Bulletin of Zoology, Supplement No. 13*, 175–208.
- Parish, F. (2002). Overview on peat, biodiversity, climate change and fire. In F. Parish, E. Padimanabhan, C.L. Lee, & H.C. Thang (eds.), *Prevention and Control of Fire in Peatlands, 19-21 March 2002*. Global Environmental Centre and Forestry Department, Peninsular Malaysia. Cetaktama, Kuala Lumpur, Malaysia.
- PIU-SERT. (2013). Studi ilmiah untuk rehabilitasi dan pengelolaan hutan gambut Rawa Tripa. In *Stakeholders Workshop of Tripa Peat Swamp Forest Management*. Banda Aceh: Unsyiah–Satgas Redd/UKP4.
- Posa, M. R. C., Wijedasa, L. S., & Corlett, R. T. (2011). Biodiversity and conservation of tropical peat swamp forests. *BioScience*, 61(1), 49–57.
- Prentice, C. & Parish, D. (1992). Conservation of peat swamp forest: A forgotten ecosystem. *Malayan Nature Journal*, 45, 128–144.
- Rainboth, W. J., Vidthayanon, C., & Mai, D. Y. (2012). *Fishes of the greater Mekong ecosystem with species list and photographic atlas*. Miscellaneous Publications, Museum of Zoology, University of Michigan.
- Renato, A. M., Benedito, S., Amaral, D., & Oyakawady, O. T. (2000). Spatial and temporal patterns of diversity and distribution of the upper Juruá river fish community (Brazilian amazon). *Environmental Biology of Fishes*, 57(1), 25–35.
- Reynolds, J. B., & Kolz, A. L. (1993). Electrofishing. In V.Z. Alexander, D. L. Parrish, & T.M. Sutton (Eds.), *Fisheries Technique*. Bethesda, Maryland: American Fisheries Society.
- Rezawaty, K. (2004). Taburan ikan di tiga batang sungai di Paya Beriah. B.Sc. thesis, Universiti Sains Malaysia, Pulau Pinang.
- Roberts, T. R. (1993). The freshwater fishes of Java, as observed by Kuhl & van Hasselt in 1820–1823. *Zoologische Verhandelingen*, 285(1), 1–94.
- Rydin, H., & Jeglum, J. K. (Eds.) (2006). *The Biology of Peatlands*. Oxford: Oxford University Press.
- Sade, A., & Biun, H. (2012). The ichthyofauna of Maliau Basin buffer zone at Maliau Basin conservation area, Sabah, Malaysia. *Journal of Tropical Biology and Conservation*, 9, 105–113.
- Shah, A. S. R. M., Zarul, H. H., Chan, K. Y., Zakaria, R., Khoo, K. H., & Mashhor, M. (2006). A recent survey of freshwater fishes of the Paya Beriah peat swamp forest, north Perak, Malaysia. *Jurnal Biosains*, 17(1), 51–64.
- Sharber, N. G. & Black, J. S. (1999). Epilepsy as a unifying principle in electrofishing theory: A proposal. *Transactions of the American Fisheries Society*, 128(4), 666–671.
- Shiraishi, Y., Mizuno, N., Nagai, M., Yoshimi, M., & Nishiyama, K. (1972). Studies on the diel activity and feeding habit of fishes at Lake Bera, Malaysia. *Japanese Journal of Ichthyology*, 19(4), 295–306.

- Siow, R., Ramli, M. N., Shakori, M. H. M., & Asmuni, M. (2013). *Fish fauna assemblages and their distribution patterns in the north Selangor peat swamp forest*. Proceedings of the Seminar on Scientific Biodiversity Expedition, North Selangor Peat Swamp Forest, 28th November 2013, Quality Hotel, Shah Alam, Selangor, Malaysia. pp.77–82.
- Tabacchi, E., Platy-Tabacchi, A. M., & Decamps, O. (1998). Continuity and discontinuity of riparian vegetation along a fluvial corridor. *Landscape Ecology*, 5(1), 9–20.
- Tan, H. H., & Kottelat, M. (2009). The fishes of Batang Hari drainage, Sumatra, with description of six new species. *Ichthyological Exploration of Freshwaters*, 20(1), 13–69.
- Tan, H. H., & Lim, K. K. P. (2013). Three new species of freshwater halfbeaks (Teleostei: Zenarchopteridae: Hemirahamphodon) from Borneo. *The Raffles Bulletin of Zoology*, 61(2), 735–747.
- Taskforce REDD. (2012). *Visi dan Misi Pemulihan Kondisi Lahan Gambut Rawa Tripa*. Initial Report Satgas REDD/UKP4, Jakarta.
- Thompson, K. G., Bergersen, E. P., & Nehring, R. B. (1997). Injuries to brown trout and rainbow trout induced by capture with pulsed direct current. *North American Journal of Fisheries Management*, 17(1), 141–153.
- Thompson, K. G., Bergersen, E. P., Nehring, R. B., & Bowden, D. C. (1997). Long-term effects of electrofishing on growth and body condition of brown trout and rainbow trout. *North American Journal of Fisheries Management*, 17(1), 154–159.
- Torres, A. G. (2000). *Labiobarbus lineatus*. *Fish Base Online*. Retrieved from www.fishbase.se/summary/60498.
- UNDP. (2006). *Malaysia's Peat Swamp Forests: Conservation and Sustainable Use. International Journal of Integrated Engineering – Special Issue on ICONCEES*, 3. United Nations Development Programme. Kuala Lumpur, Malaysia.
- Van der Werf, G. R., Morton, D. C., DeFries, R. S., Olivier, J. G. J., Kasibhatla, P. S., Jackson, R. B., Collatz, G. J., & Randerson, J. T. (2009). CO₂ emissions from forest loss. *Nature Geoscience*, 2(11), 737–739.
- Wantzen, K. M., Yule, C. M., Mathooko, J. M., & Pringle, C. M. (2011). Organic matter processing in tropical streams. *Tropical Stream Ecology*, 1, 44–65.
- Welcomme, R. L. (1979). *Fisheries Ecology of Floodplain Rivers*. London: Longman.
- Whitmore, T. C. (1984). *Tropical Rain Forests of the Far East*. Clarendon Press, Oxford.
- Whitmore, T. C. (1988). Forest types and forest zonation. In Earl of Cranbrook (ed.), *Key Environments: Malaysia*. Elsevier.
- Wright, J. P., & Flecker, A. A. (2004). Deforesting the riverscape: The effects of food on fish diversity in a Venezuelan piedmont stream. *Biological Conservation*, 120(3), 443–451.
- Wyatt-Smith, J. (1959). Peat swamp forests in Malaya. *Forester*, 22(1), 5–31.
- Wyatt-Smith, J. (1964). A preliminary vegetation map of Malaya with description of the vegetation types. *Journal of Tropical Geography*, 18, 200–213.
- Yule, C. M. (2010). Loss of biodiversity and ecosystem functioning in Indo-Malayan peat swamp forests. *Biodiversity and Conservation*, 19(2), 393–409.
- Zakaria, R., Mansor, M., & Ali, A. B. (1999). Swamp-riverine tropical fish population : A comparative study of two spatially isolated freshwater ecosystems in Peninsular Malaysia. *Wetlands Ecology and Management*, 6(4), 261–268.
- Zakaria-Ismail, M. (1990). Two-spotted catfish. *Nature Malaysiana*, 15(3), 88–89.